

brexanolone	<ul style="list-style-type: none"> ❗ Restricted to [REDACTED]
brexpiprazole	<ul style="list-style-type: none"> ❗ restricted to continuation of home meds
Bupivacaine and Meloxicam	<ul style="list-style-type: none"> ❗ Restricted to [REDACTED] and [REDACTED] Hospitals only. Restricted to open mesh inguinal and ventral hernias.
bupivacaine liposome	<ul style="list-style-type: none"> ❗ Transversus abdominis plane (TAP) block for all open abdominal surgery ❗ Interscalene block for major shoulder surgery ❗ Current/active IRB-approved research studies ❗ All surgical cases involving Epidermolysis Bullosa (EB) patients that meet the FDA-approved age recommendations
Buprenorphine	<ul style="list-style-type: none"> ❗ (Brixadi Monthly) Outpatient infusion centers and clinics ❗ (Brixadi Weekly) Pilot program through [REDACTED] Emergency Department (ED) for adult or pediatric patients (note this does not include use in the [REDACTED] Pediatric ED or the BEC) ❗ (Brixadi Weekly) For patients that may be admitted, discussion between the ED provider and admitting team should occur before administration of Brixadi in the [REDACTED] ED; outpatient infusion centers and clinics.
bupropion-dextromethorphan	<ul style="list-style-type: none"> ❗ Outpatient, continuation of home medication, restrict to psychiatry
cabotegravir-rilpivirine	<ul style="list-style-type: none"> ❗ If an established patient is admitted inpatient, and the patient will miss their outpatient injection by > 7 days, Cabenuva may be ordered and administered while an inpatient.
cabozantinib	<ul style="list-style-type: none"> ❗ Use patient's own supply
calaspargase pegol	<ul style="list-style-type: none"> ❗ Restricted to clinic and inpatient for induction.
cannabidiol	<ul style="list-style-type: none"> ❗ Epidiolex: Patients may use their own supply