

Restricted

Please review each of these drugs carefully.

abatacept	<p>i abatacept (Orencia) is restricted to use in the Outpatient setting by Rheumatology only</p>
acetaminophen	<p>i Acetaminophen IV (Ofirmev) - See comments for restrictions</p>
acetylcysteine	<p>i acetylcysteine (Acetadote/Mucomyst) nebulization - use by intensivists (Trauma, Med/Surge, ICU) in patients who failed nebulization of 3% NaCl</p> <p>i acetylcysteine (Acetadote/Mucomyst) oral and IV - is restricted to treatment of APAP toxicity and non-acetaminophen induced acute liver injury. All other indications must be approved.</p>
acyclovir topical	<p>i formulary restricted as alternative to docosanol when application to large surface area is required</p>
adalimumab	<p>i Restricted to outpatient use only</p>
ado-trastuzumab emtansine	<p>i ado-trastuzumab emtansine (Kadcyla) is restricted to use in the Outpatient setting</p>
alemtuzumab	<p>i alemtuzumab (Lemtrada) is restricted to use in the Outpatient setting and is a REMS medication</p>
alglucosidase alfa	<p>i Restricted to outpatient setting only</p>
aminolevulinic acid	<p>i Aminolevulinicacid (Gleolan) will be formulary restricted to surgeons who have completed appropriate training and have appropriate</p>