

Legacy Health DTU Nonhazardous IV Admixture Standards

Drug	Route: Standard Concentration and Diluent	End-product BUD/Hang by	Special Instructions	Attachments: Tubing, Filter	Aux Labels	Vial Size and Storage Fridge unless noted	Vial Reconstitution	Vial Exp. Fridge unless noted	Reference I=Insert, T=Trissels, M=Micromedex, A=Article, F&C= Facts & Comparisons
abatacept (Orencia)	IV: no more than 10 mg/mL in total volume 100 mL NS	24 h	Use silicone free syringe Aim SWFI at vial wall, Do not shake. May take up to 30 min to dissolve. Total volume = 100mL (withdraw a volume equal to the dose from the bag, before adding the dose)	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	250 mg	10 mL SWFI	discard unused	I (ER Squibb, 11/2025)
agalsidase beta (Fabrazyme)	IV: 0.2 - 0.7 mg/mL in total volume 100 mL or 250 mL NS	4 h	Bring vials to room temp, ~30 min before reconstitution; Do NOT shake Do NOT inject into air space of IVPB If not administered immediately, may store 24 h FRIDGE	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	5 mg 35 mg	5mg = 1.1mL SWFI 35mg = 7.2mL SWFI	discard unused	I (Genzyme/Sanofi, 8/2025)
alpha-1 proteinase inhibitor (Prolastin-C)	IV: Dosed at 60 mg/kg Do NOT dilute	3 h	Bring vials to room temperature Transfer into empty IV bag Administer within 3 hours preparation	5 micron filter (474002)	Filter	~500 mg/10 mL ~1000 mg/20 mL ~4000 mg/80 mL	n/a	discard unused	I (Grifols, 7/2022)
amivantamab-vmjw (Rybrevant)	IV: Dose in total volume 250 mL NS or D5W	10 h	Total volume = 250mL (withdraw amount of fluid from bag equal to dose being added, before adding the dose). Do NOT shake.	Send with NS or D5W PRIMED Low sorbing tubing + 0.2 micron filter (11532269)	Filter	350 mg/7 mL	n/a	discard unused	I (Janssen, 11/2025)
aprepitant (Cinvanti)	IV: Dose in 100 mL NS or D5W	6 h (NS) 12 h (D5W)	Gently invert bag to mix. Avoid shaking.			130 mg/18 mL	n/a	discard unused	I (Heron, 3/2024)
atezolizumab (Tecentriq)	IV: 3.2 - 16.8 mg/mL NS	6 h	Gently invert bag to mix. Do not shake. If not administered immediately, may store 24 h FRIDGE			840 mg/14 mL 1200 mg/20 mL	n/a	discard unused	I (Genetech, 11/2025)
avelumab (Bavencio)	IV: Dose in 250mL NS	4 h	If not administered immediately, may store 24 h FRIDGE	LPB 0.2 micron filter (20027E)	Protect from Light Filter	200 mg/10 mL	n/a	discard unused	I (EMD Serono, 6/2025)
belatacept (Nulojix)	IV: 2 - 10 mg/mL total volume NS or D5W	4 h	Use silicone free syringe; direct stream of fluid to side wall; do not shake. Withdraw amount of fluid from bag equal to dose being added, before adding the dose If not administered immediately, may store 24 h FRIDGE	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	250 mg	10.5 mL SWFI, NS, D5W	discard unused	I (ER Squibb, 7/2021)
belimumab (Benlysta)	IV: NTE 4 mg/mL total volume in 100 mL or 250 mL NS	8 h	Use 21 to 25ga needle; Direct SWFI at wall of vial; Gently swirl for 60 sec every 5 min, up to 30 min; Withdraw amount of fluid from bag equal to dose being added, before adding the dose			120 mg 400 mg	120 mg = 1.5 mL SWFI 400 mg = 4.8 mL SWFI May take up to 30 min to dissolve DO NOT SHAKE	discard unused	I (GSK, 6/2025)
bevacizumab (Avastin)	IV: Dose in 100 mL total volume NS	discard by 8 h Fridge	NS ONLY		Refrigerate	100 mg/4 mL 400 mg/16 mL	n/a	discard unused	I (Genetech, 1/2025)
bevacizumab-awwb (Mvasi)	IV: Dose in 100 mL total volume NS	discard by 8 h Fridge	NS ONLY		Refrigerate	100 mg/4 mL 400 mg/16 mL	n/a	discard unused	I (Amgen, 10/2024)
bevacizumab-bvzr (Zirabev)	IV: Dose in 100 mL total volume NS	8 h Fridge	NS ONLY If not administered immediately, may store 16 d FRIDGE		Refrigerate	100 mg/4 mL 400 mg/16 mL	n/a	discard unused	I (Pfizer, 8/2025)

NOTE: BUD times reflect admixture in an ISO 5 LAFW within an ISO 7 environment. SCA compounding reduces BUD to 12 h room temp, 24 h fridge unless stability of product is less.

Approved by USP Standards, 2/2026

Legacy Health DTU Nonhazardous IV Admixture Standards

Drug	Route: Standard Concentration and Diluent	End-product BUD/Hang by	Special Instructions	Attachments: Tubing, Filter	Aux Labels	Vial Size and Storage Fridge unless noted	Vial Reconstitution	Vial Exp. Fridge unless noted	Reference I=Insert, T=Trissels, M=Micromedex, A=Article, F&C= Facts & Comparisons
carfilzomib (Kyprolis)	IV: Dose in 50 or 100 mL D5W	4 h	Gently invert bag to mix. If not administered immediately, may store 24 h FRIDGE			10 mg 30 mg 60 mg	10 mg = 5 mL SWFI 30 mg = 15 mL SWFI 60 mg = 29 mL SWFI	discard unused	I (Onyx Pharmaceuticals, 6/2022)
cemiplimumab (Libtayo)	IV: 1 mg/mL to 20 mg/mL in NS or D5W	8 h	Gently invert bag to mix. If not administered immediately, may store 10 d FRIDGE	LPB 0.2 micron filter (20027E)	Filter	350 mg/7 mL	n/a	discard unused	I (Regeneron, 10/2025)
cetuximab (Erbix)	IV syringe: 2 mg/mL Do NOT dilute	8 h	Do not administer IV push or bolus	LPB 0.2 micron filter (20027E)	Filter	100 mg/50 mL 200 mg/100 mL	n/a	8 h RT 12 h Fridge	I (ImClone, 7/2024)
crizanlizumab (Adakveo)	IV: Dose in total volume 100 mL NS or D5W	4.5 h	Bring vials to RT before prep (up to 4 h). Total volume = 100mL (withdraw volume equal to dose from bag, before adding). Gently invert bag to mix. Do not shake. If not administered immediately, may store 24 h FRIDGE	LPB 0.2 micron filter (20027E)	Filter	100 mg/10 mL	n/a	discard unused	I (Novartis, 6/2025)
dalbavancin (Dalvance)	IV: 1-5 mg/mL in D5W	48 h	Gently swirl, alternate with inverting vial to avoid foaming. Do not shake. Dilute in D5W only			500 mg/25 mL Room Temp	25 mL SWFI or D5W	discard unused	I (Allergan, 1/2025)
daratumumab (Darzalex)	IV: Dose in total volume 500 mL or 1000 mL NS	15 h	Gently invert bag to mix. Do not shake. If not administered immediately, may store 24 h FRIDGE	LPB 0.2 micron filter (20027E)	Filter	100 mg/5 mL 400 mg/20 mL	n/a	discard unused	I (Janssen, 4/2025)
daratumumab + hyaluronidase-fihj (Darzalex Faspro)	SC: Do NOT dilute	12 h	Remove vial from refrigerated storage & equilibrate to ambient temperature. If not administered immediately, may store 24 h FRIDGE			1800 mg + 30,000 units/ 15mL	n/a	discard unused	I (Janssen, 12/2025)
degarelix (Firmagon)	SC: 40 mg/mL	1 h	240 mg dose is composed of 2 x 120 mg doses			80 mg powder kit 120 mg powd kit Room Temp	80 mg:4.2mL SW in kit 120 mg:3 mL SW in kit	discard unused	I (Ferring Pharm, 1/2024)
dinutuximab (Unituxin)	IV: Dose in 100 mL NS	Discard by 4 h Fridge	Gently invert bag to mix. Do not shake. If not administered immediately, may store 24 h FRIDGE			17.5 mg/5 mL	n/a	discard unused	I (United Therapeutics, 10/2025)
dostarlimab-gxly (Jemperli)	IV: Dose in 250 mL NS or D5W 500 mg dose: final concentration, 2-10 mg/mL 1000 mg dose: final concentration, 4-10 mg/mL	6 h	If not administered immediately, may store 24 h FRIDGE	LPB 0.2 micron filter (20027E)	Filter	500 mg/10 mL	n/a	discard unused	I (GlaxoSmithKline, 9/2025)
durvalumab (Imfinzi)	IV: 1 - 15 mg/mL in NS or D5W	8 h	Gently invert bag to mix. Do not shake. If not administered immediately, may store 28 d FRIDGE	LPB 0.2 micron filter (20027E)	Filter	120 mg/2.4 mL 500 mg/10 mL	n/a	discard unused	I (AstraZeneca, 11/2025)
eculizumab (Soliris)	IV: 5 mg/mL in NS or D5W	24 h	Use equal amounts of drug and diluent in empty bag. Gently invert to mix.			300 mg/30 mL	n/a	discard unused	I (Alexion, 3/2025)
eculizumab-aagh (Epysqli)	IV: 5 mg/mL NS or D5W	24 h	Use equal amounts of drug and diluent in empty bag. Gently invert to mix.			300 mg/30 mL	n/a	discard unused	I (Teva, 12/2025)
efgartigimod alfa-fcab (Vyvgart)	IV: Dose in total volume 125 mL NS	4 h 8 h Fridge	Gently invert to mix. Do not shake.	LPB 0.2 micron filter (20027E)	Filter	400 mg/20 mL	n/a	discard unused	I (Argenx, 10/2025)
elosulfase alfa (Vimizim)	IV: <25 kg: total vol 100 mL NS >=25 kg: total volume 250 mL NS	24 h	Gently invert to mix. Do not shake.	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	5 mg/5 mL	n/a	discard unused	I (Biomarin, 10/2025)

NOTE: BUD times reflect admixture in an ISO 5 LAFW within an ISO 7 environment. SCA compounding reduces BUD to 12 h room temp, 24 h fridge unless stability of product is less.

Approved by USP Standards, 2/2026

Legacy Health DTU Nonhazardous IV Admixture Standards

Drug	Route: Standard Concentration and Diluent	End-product BUD/Hang by	Special Instructions	Attachments: Tubing, Filter	Aux Labels	Vial Size and Storage Fridge unless noted	Vial Reconstitution	Vial Exp. Fridge unless noted	Reference I=Insert, T=Trissels, M=Micromedex, A=Article, F&C= Facts & Comparisons
elotuzumab (Empliciti)	IV: 1 to 6 mg/mL in NS or D5W	8 h	Use less than a 18ga needle. Gently swirl. Allow to stand for 5 to 10 min. If not administered immediately, may store 24 h FRIDGE	LPB 0.2 micron filter (20027E)	Filter	300 mg 400 mg	300 mg = 13 mL SWFI 400 mg = 17 mL SWFI	discard unused	I (ER Squibb, 3/2022)
elranatamab-bcmm (Elrexfio)	SC: Do NOT dilute	24 h	Remove vial from refrigerated storage and equilibrate to ambient temperature. If not administered immediately, may store 72 h FRIDGE			44 mg/1.1 mL 76 mg/1.9 mL	n/a	discard unused	I (Pfizer, 7/2025)
eptinezumab-jjmr (Vyepiti)	IV: Dose in total volume 100 mL NS	8 h	Gently invert to mix. Do not shake.	LPB 0.2 micron filter (20027E)	Filter	100 mg/1 mL	n/a	discard unused	I (Lundbeck, 10/2025)
epcoritamab-bysp (Epkinyly)	SC: 3 mg and 48 mg doses, Do NOT dilute. 0.16 mg and 0.8 mg doses, dilute with NS.	12 h	During preparation allow the vial to come to room temp for no more than 1 hr. If not administered immediately, may store 24 h FRIDGE			4 mg/0.8 mL 48 mg/0.8 mL	n/a	discard unused	I (Genmab, 11/2025)
ferric carboxymatlose (Injectafer)	IV: not less than 2 mg/mL NS	72 h				100 mg/2 mL 750 mg/15 mL 1000 mg/20 mL Room Temp	n/a	discard unused	I (American Regent, 1/2025)
ferric gluconate (Ferrelecit)	125 mg in 110 mL NS 250 mg in 270 mL NS	24 h	Use immediately			62.5 mg/5 mL Room Temp	n/a	discard unused	I (Hikma, 7/2022) A (Baribeault, 2011)
ferumoxytol (Feraheme)	IV: 2-8 mg/mL elemental iron in 50-200 mL NS or D5W	4 h	If not administered immediately, may store 48 h FRIDGE			510 mg/17 mL Room Temp	n/a	discard unused	I (AMAG, 6/2022)
fosaprepitant (Emend)	IV: 150 mg in total volume 250 mL NS	24 h	Do not mix in LR. Swirl vial gently. Gently invert bag to mix. Do not shake.			150 mg	5 mL NS Add along vial wall to prevent foaming, swirl	discard unused	M, I (Hikma, 10/2021)
golimumab (Simponi Aria)	IV: Dose in total volume 100 mL NS	4 h	Gently invert to mix. Do not shake.	LPB 0.2 micron filter (20027E)	Filter	50 mg/4 mL	n/a	discard unused	I (Janssen, 9/2025)
guselkumab (Tremfya)	IV: 0.8 mg/mL in total volume 250 mL NS	10 h	Gently invert to mix. Do not shake.	LPB 0.2 micron filter (20027E)	Filter	200 mg/20 mL	n/a	discard unused	I (Janssen, 10/2025)
imiglucerase (Cerezyme)	IV: Dose in 100 to 200 mL NS	24 h Fridge	When reconstituting, slowly inject SWFI down inside wall of vial. Roll and tilt to dissolve.	LPB 0.2 micron filter (20027E)	Filter	400 units	400 units = 10.2 mL SWFI	12 h RT or Fridge	I (Genzyme, 8/2025)
immune globulin	IV: straight draw into empty bag - 10% solution	8 h	Do not use if the solution is cloudy, turbid or contains particulate matter. Do not shake.			5 g/50 mL 10 g/100 mL 20 g/200 mL 40 g/100 mL Room Temp	n/a	discard unused	I (CSL Behring, 5/2025)
infliximab (Remicade)	IV: 0.4 - 4 mg/mL in total volume 250 mL NS	3 h	Gently swirl to dissolve powder. Allow the reconstituted solution to stand for 5 min. Gently invert bag to mix. Do not shake.	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	100 mg	100 mg = 10 mL SWFI	discard unused	I (Janssen, 6/2025)
infliximab-axxq (Avsola)	IV: 0.4 - 4 mg/mL in total volume 250 mL NS	3 h	Gently swirl to dissolve powder. Allow the reconstituted solution to stand for 5 min. Gently invert bag to mix. Do not shake.	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	100 mg	100 mg = 10 mL SWFI	discard unused	I (Amgen, 9/2025)
infliximab-dyyb (Inflectra)	IV: 0.4 - 4 mg/mL NS Doses > 1000 mg must be diluted in 500 mL NS	3 h	Gently swirl to dissolve powder. Allow the reconstituted solution to stand for 5 min. Gently invert bag to mix. Do not shake.	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	100 mg	100 mg = 10 mL SWFI	discard unused	I (Pfizer, 9/2025), F&C

NOTE: BUD times reflect admixture in an ISO 5 LAFW within an ISO 7 environment. SCA compounding reduces BUD to 12 h room temp, 24 h fridge unless stability of product is less.

Approved by USP Standards, 2/2026

Legacy Health DTU Nonhazardous IV Admixture Standards

Drug	Route: Standard Concentration and Diluent	End-product BUD/Hang by	Special Instructions	Attachments: Tubing, Filter	Aux Labels	Vial Size and Storage Fridge unless noted	Vial Reconstitution	Vial Exp. Fridge unless noted	Reference I=Insert, T=Trissels, M=Micromedex, A=Article, F&C= Facts & Comparisons
ipilimumab (Yervoy)	IV: 1 - 2 mg/mL NS or D5W	24 h	Allow to warm to room temp for 5 min prior to preparation.	LPB 0.2 micron filter (20027E)	Filter	50 mg/10 mL 200 mg/40 mL	n/a	discard unused	I (ER Squibb, 5/2025)
iron dextran	IV: 2 - 4 mg/mL, 500 mg or 1000 mg in 250 mL NS	4 d	Do not remove overflow or drug volume from diluent bag			100 mg/2 mL Room Temp	n/a	6 h	USP, I (Allergan, 8/2024)
iron sucrose	IV: concentration ≥ 1 mg 100 - 200 mg in 100 mL NS > 200 mg in 250 mL NS	4 d				100 mg/5 mL Room Temp	n/a	6 h	USP, I (American Regent, 6/2022)
isatuximab-irfc (Sarclisa)	IV: Dose in total volume 250 mL NS or D5W	8 h	Gently invert to mix. Do not shake. If not administered immediately, may store 48 h FRIDGE	LPB 0.2 micron filter (20027E)	Filter	100 mg/5 mL 500 mg/25 mL	n/a	discard unused	I (Sanofi-Aventis, 6/2025)
leucovorin	IVPB: in 250 mL NS or D5W Syringe: 10 mg/mL SWFI	24 h	Do NOT reconstitute using bacteriostatic water		Protect from light	50 mg, 100 mg, 200 mg, 350 mg Room Temp	50 mg: 5 mL SW 100 mg: 10 mL SW 200 mg: 20 mL SW 350 mg: 17.5 mL SW	discard unused	T, I (Sagent, 2020; Mylan, 2022; Fresenius Kabi, 3/2022; Methial, 5/2024) A (Benevenuto, accessed 2025)
luspatercept-aamt (Reblozyl)	SC: 50 mg/mL SWFI	8 h	Step 1: Gently swirl reconstituted vial for 30 sec, rest for 30 s, & repeat until dissolved. Step 2: invert the vial & swirl for 30 s, rest for 30 s, repeat step 7 times. If not administered immediately, may store 24 h FRIDGE			25 mg 75 mg	25 mg=0.68 mL SWFI 75 mg=1.6 mL SWFI	discard unused	I (Celgene, 5/2024)
margetuximab-cmkb (Margenza)	IV: 0.5 to 7.2 mg/mL in 100 or 250 mL NS	4 h	Gently invert bag to mix. Do not shake. If not administered immediately, may store 24 h FRIDGE	LPB 0.2 micron filter (20027E)	Filter	250 mg/10 mL	n/a	discard unused	I (TerSera, 2/2025)
mesna (Mesnex)	Up to 20 mg/mL in NS or D5W	24 h				1000 mg/10 mL Room Temp	n/a	8 d	F&C, I (Sagent, 3/2025)
methylprednisolone	IV: 1-20 mg/mL NS or D5W	48 h	Doses of 125 mg or less: use of Act-O-Vial preferred via IV push		Protect from Light	Act-O-Vial: 40 mg, 125 mg 500 mg, 1 g powder Room Temp	500 mg: 8 mL Bact SW 1 g: 16 mL Bact SW	48 h	I (Pfizer, 3/2024), T, A (Trissel, 2002)
mogamulizumab-kpkc (Poteligeo)	IV: 0.1 to 3 mg/mL NS	24 h Fridge	Gently invert bag to mix. Do not shake.	LPB 0.2 micron filter (20027E)	Filter	20 mg/5 mL	n/a	discard unused	I (Kyowa Kirin, 10/2025)
mosunetuxumab-axgb (Lunsumio)	IV: Dose in total volume 50 mL, 100 mL, or 250 mL NS	16 h RT, 24 h Fridge	Withdraw amount of fluid from bag equal to dose being added, before adding the dose Gently invert to mix. Do not shake.			1 mg/mL 30 mg/30 mL	n/a	discard unused	I (Genetech, 12/2025)
multivitamin-containing IVPB (aka Banana Bag)	IV: in 500 mL NS or D5W	24 h	Once MVI vials are combined with IVPB, discard bag by 24 h			2 vials, 5 mL ea	transfer contents of vial 1 into the contents of vial 2	discard unused	I (Baxter, 11/2024)
nadofaragene firadenovec-vncg (Adstiladrin)	Intravesical admin: Do NOT dilute	6 h	Bring to room temperature prior to use. Time for thawing/bringing vials to room temp is ~8 to 10 h when thawing in the cardboard nest, and ~3 to 5 h when thawing vials outside the cardboard nest.		Protect from Light	3 × 10 ¹¹ viral particles (vp)/mL in 20 mL Freezer	n/a	discard unused	I (Ferring, 10/2025)
natalizumab (Tysabri)	IV: 2.6 mg/mL in 100 mL NS	48 h Fridge	Gently invert bag to mix. Do not shake.		Refrigerate	300 mg/15 mL	n/a	discard unused	T, I (Biogen, 3/2025)

NOTE: BUD times reflect admixture in an ISO 5 LAFW within an ISO 7 environment. SCA compounding reduces BUD to 12 h room temp, 24 h fridge unless stability of product is less.

Approved by USP Standards, 2/2026

Legacy Health DTU Nonhazardous IV Admixture Standards

Drug	Route: Standard Concentration and Diluent	End-product BUD/Hang by	Special Instructions	Attachments: Tubing, Filter	Aux Labels	Vial Size and Storage Fridge unless noted	Vial Reconstitution	Vial Exp. Fridge unless noted	Reference I=Insert, T=Trissels, M=Micromedex, A=Article, F&C= Facts & Comparisons
naxitamab-gqgk (Danyelza)	IV: Dose in 5% Albumin + NS	8 h	Add albumin and NS to empty IV bag, allow 5 to 10 min of passive mixing. Inject drug into albumin/NS containing bag.			40 mg/10 mL	n/a	discard unused	I (Y-mAbs, 9/2025)
nivolumab (Opdivo)	IV: 1 - 10 mg/mL NS or D5W	8 h	Total volume not to exceed 160 mL. Gently invert bag to mix. Do not shake. If not administered immediately, may store 7 d FRIDGE,	LPB 0.2 micron filter (20027E)	Filter	40 mg/4 mL 100 mg/10 mL 120 mg/12 mL 240 mg/24 mL	n/a	discard unused	I (ER Squibb, 7/2025)
nivolumab and relatlimab (Opdualag)	IV: Nivolumab: 3 - 12 mg/mL Relatlimab: 1 - 4 mg/mL in NS or D5W	8 h	May administer undiluted or diluted with a total volume not to exceed 160 mL. Gently invert bag to mix. Do not shake. If not administered immediately, may store 24 h FRIDGE	LPB 0.2 micron filter (20027E)	Filter	240 mg & 80 mg/20 mL	n/a	discard unused	I (ER Squibb, 3/2024)
obintuzumab (Gazyva)	IV: 0.4 - 4 mg/mL NS	24 h Fridge	Cycle 1 doses, day 1 (100 mg) and day 2 (900 mg) prepared on same day in buffer room. Store day 2 dose x 24 h fridge.			1000 mg/40 mL	n/a	discard unused	I (Genetech, 12/2025)
ocrelizumab (Ocrevus)	IV: 1.2 mg/mL in 250 mL or 500 mL NS	8 h	If not administered immediately, may store 24 h FRIDGE	LPB 0.2 micron filter (20027E)	Filter	300 mg/10 mL	n/a	discard unused	I (Genetech, 12/2025)
omalizumab (Xolair)	SC: 125 mg/mL SWFI	4 h	Use 1" 18 ga needle. Gently swirl vial for 5 to 10 seconds every 5 minutes. Takes 15 to 20 min to dissolve.			150 mg	150 mg: 1.4 mL SWFI	4 h RT 8 h Fridge	I (Genetech, 12/2025)
panitumumab (Vectibix)	IV: NTE 10 mg/mL in total volume 100 mL NS Doses > 1g in 150 mL NS	6 h	Do not use spikes on vials. Gently invert bag to mix. Do not shake. If not administered immediately, may store 24 h FRIDGE	LPB 0.2 micron filter (20027E)	Filter	100 mg/5 mL 400 mg/20 mL	n/a	discard unused	I (Amgen, 6/2025)
pegloticase (Krystexxa)	IV: Dose in 250 mL NS	4 h	Gently invert bag to mix. Do not shake.			8 mg/1 mL	n/a	discard unused	I (Horizon, 8/2025)
pembrolizumab (Keytruda)	IV: 1 - 10 mg/mL in NS or D5W	6 h	Gently invert bag to mix. Do not shake. If not administered immediately, may store 96 h FRIDGE	LPB 0.2 micron filter (20027E)	Filter	100 mg/4 mL	n/a	discard unused	I (Merck Sharp & Dohme, 11/2025)
pemivibart (Pemgarda)	IV: Dose in total volume 50 mL NS	4 h	Remove vials from refrigerator at least 10 minutes before preparation. Gently invert bag to mix. Do not shake.	LPB 0.2 micron filter (20027E)	Filter	500 mg/4 mL	n/a	discard unused	I (Inivyd, 10/2025)
pentamidine	IV: Dose in 50-250 mL D5W	24 h				300 mg Room Temp	300 mg: 3-5 mL SWFI	48 h SCA: 12 h Protect from Light	T, I (Seton, 4/2025)
pertuzumab (Perjeta)	IV: Dose in 250 mL NS	24 h Fridge	Gently invert bag to mix. Do not shake.			420 mg/14 mL	n/a	discard unused	I (Genentech, 12/2025)
pertuzumab/ trastuzumab and hyaluronidase-zzxf (Phesgo)	SC: 10 mL or 15 mL Do NOT dilute	4 h	Do not split syringes - send full volume in one syringe. If not administered immediately, may cap syringe and store 24 h Fridge			1200/600/30,000 /15 mL 600/600/20,000 /10 mL	n/a	discard unused	I (Genentech, 12/2025)
ramucirumab (Cyramza)	IV: Dose in total volume 250 mL NS	4 h	If not administered immediately, may store 24 h Fridge	LPB 0.2 micron filter (20027E)	Filter	100 mg/10 mL 500 mg/50 mL	n/a	discard unused	I (Eli Lilly, 12/2025)
ravulizumab-cwvz (Ultomiris)	IV: 50 mg/mL in NS	4 h	Gently invert bag to mix. Do not shake. If not administered immediately, may store 24h Fridge	LPB 0.2 micron filter (20027E)	Protect from Light Filter	300 mg/3 mL 1100 mg/11 mL	n/a	discard unused	I (Alexion, 10/2025)

NOTE: BUD times reflect admixture in an ISO 5 LAFW within an ISO 7 environment. SCA compounding reduces BUD to 12 h room temp, 24 h fridge unless stability of product is less.

Approved by USP Standards, 2/2026

Legacy Health DTU Nonhazardous IV Admixture Standards

Drug	Route: Standard Concentration and Diluent	End-product BUD/Hang by	Special Instructions	Attachments: Tubing, Filter	Aux Labels	Vial Size and Storage Fridge unless noted	Vial Reconstitution	Vial Exp. Fridge unless noted	Reference I=Insert, T=Trissels, M=Micromedex, A=Article, F&C= Facts & Comparisons
reslizumab (Cinqair)	IV: Dose in 50 mL NS	16 h	Gently invert bag to mix. Do not shake.	LPB 0.2 micron filter (20027E)	Filter	100 mg/10 mL	n/a	discard unused	I (Teva, 11/2025)
retifanlimab-dlwr (Zynyz)	IV: 1.4 - 10 mg/mL in NS or D5W	8 h	Gently invert bag to mix. Do not shake. If not administered immediately, may store 24 h Fridge	LPB 0.2 micron filter (20027E)	Filter	500 mg/20 mL	n/a	discard unused	I (Incyte, 12/2025)
risankizumab-rzaa (Skyrizi)	IV: 1.2 - 6 mg/mL in NS or D5W	4 h	Gently invert bag to mix. Do not shake. If not administered immediately, may store 20 h Fridge Exposure to indoor light during RT storage and administration is acceptable.		Protect from Sunlight	600 mg/10 mL	n/a	discard unused	I (Abbvie, 9/2025)
rituximab (Rituxan)	IV: 1 - 4 mg/mL NS or D5W	24 h	Gently invert bag to mix. Do not shake.			100 mg/10 mL 500 mg/50 mL	n/a	discard unused	I (Genetech, 1/2025)
rituximab-abbs (Truxima)	IV: 1 - 4 mg/mL NS or D5W	24 h Fridge	Gently invert bag to mix. Do not shake.			100 mg/10 mL 500 mg/50 mL	n/a	discard unused	I (Teva/Cephalon, 6/2025)
rituximab-pvvr (Ruxience)	IV: 1 - 4 mg/mL NS or D5W	4 d Fridge (NS)	Gently invert bag to mix. Do not shake. If mixed in D5W , 24 h Fridge storage after preparation			100 mg/10 mL 500 mg/50 mL	n/a	discard unused	I (Pfizer, 6/2025)
rituximab/hyaluronidase human (RITUXAN HYCELA)	SC: Do NOT dilute	8 h	Draw up using 20 mL syringe and 25 to 30 gauge needle If not used immediately, may store 48 h Fridge			1400mg/11.7 mL 1600mg/13.4 mL	n/a	discard unused	I (Genetech, 12/2025)
romiplostim (Nplate)	SC: Doses less than 23 mcg require further dilution with NS.	4 h	Gently swirl and invert vial to reconstitute. Do not shake. Use a 1 mL syringe, or syringe with graduations to 0.01 mL. If not used immediately, may store 24 h Fridge		Protect From Light	125 mcg 250 mcg 500 mcg	125 mcg: 0.44 mL SW 250 mcg: 0.72 mL SW 500 mcg: 1.2 mL SW	discard unused	I (Amgen, 3/2025)
siltuximab (Sylvant)	IV: Dose in total volume 250 mL D5W	4 h	Gently swirl reconstituted vial. Do not shake. Further dilute within 2 h. Gently invert bag to mix.	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	100 mg 400 mg	100 mg: 5.2 mL SWFI 400 mg: 20 mL SWFI	discard unused	I (Recordati, 8/2024)
sodium ferric gluconate complex (Ferrlecit)	See Ferric Glucoconate								
tafasitamab-cxix (Monjuvi)	IV: 2 - 8 mg/mL in total volume 250 mL NS	12 h	Gently swirl vial to reconstitute; complete dissolution may take up to 5 min. Gently invert bag to mix. Do not shake. If not used immediately, may store 18 h Fridge			200 mg	200 mg: 5 mL SWFI	12 h RT or Fridge	I (Incyte, 6/2025)
talquetamab-tgvs (Talvey)	SC: Do NOT dilute	24 h	Bring vials to room temp for at least 15 min. Gently swirl vial for approximately 10 sec to mix. Do not shake. If dose > 2 mL, divide equally into multiple syringes.			3 mg/1.5 mL 40 mg/mL	n/a	discard unused	I (Janssen, 10/2025)
tebentafusp-tebn (Kimmtrak)	IV: Dose in 250 mcg/mL Albumin + 100 mL NS	4 h	Add albumin to NS bag, gently homogenize solution - do not shake. Inject drug into albumin/NS bag. If not used immediately, may store 24 h Fridge	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	100 mcg/0.5 mL	n/a	discard unused	I (Immunocore, 5/2025)

NOTE: BUD times reflect admixture in an ISO 5 LAFW within an ISO 7 environment. SCA compounding reduces BUD to 12 h room temp, 24 h fridge unless stability of product is less.

Approved by USP Standards, 2/2026

Legacy Health DTU Nonhazardous IV Admixture Standards

Drug	Route: Standard Concentration and Diluent	End-product BUD/Hang by	Special Instructions	Attachments: Tubing, Filter	Aux Labels	Vial Size and Storage Fridge unless noted	Vial Reconstitution	Vial Exp. Fridge unless noted	Reference I=Insert, T=Trissels, M=Micromedex, A=Article, F&C=Facts & Comparisons
teclistamab-cqyv (Tecvayli)	SC: Do NOT dilute	20 h	Bring vials to room temp for at least 15 min. Gently swirl vial for approximately 10 sec to mix. Do not shake. NOTE CONCENTRATION: do not combine vials of different concentrations to achieve dose. If dose > 2 mL, divide equally into multiple syringes.			30 mg/3 mL 153 mg/ 1.7 mL	n/a	discard unused	I (Janssen, 8/2025)
tocilizumab (Actemra)	IV: wt = <30 kg dose in total volume 50 mL NS wt = >30 kg dose in total volume 100 mL NS	24 h	Withdraw amount of fluid from bag equal to dose being added, before adding dose. Gently invert bag to mix. Do not shake.		Protect from Light	80 mg/4 mL 200 mg/10 mL 400 mg/20 mL	n/a	discard unused	I (Genetech, 12/2025)
tocilizumab-aazg (Tyenne)	IV: wt = <30 kg dose in total volume 50 mL NS wt = >30 kg dose in total volume 100 mL NS	24 h	Withdraw amount of fluid from bag equal to dose being added, before adding dose. Gently invert bag to mix. Do not shake.		Protect from Light	80 mg/4 mL 200 mg/10 mL 400 mg/20 mL	n/a	discard unused	I (Fresenius Kabi, 8/2025)
trastuzumab (Herceptin)	IV: Dose in 250 mL NS	24 h Fridge	Direct diluent into cake to reconstitute; swirl vial gently, allow to sit for 5 min. Gently invert bag to mix. Do not shake.			150 mg	150 mg: 7.4 mL SWFI	24 h	I (Genetech, 11/2024)
trastuzumab-anns (Kanjinti)	IV: Dose in 250 mL NS	24 h Fridge	Direct diluent into cake to reconstitute; swirl vial gently, allow to sit for 5 min. Gently invert bag to mix. Do not shake.			150 mg 420 mg	150 mg: 7.4 mL SWFI 420 mg: 20 mL BACT SW	150 mg=24 h 420 mg=28 d (w/Bact SW)	I (Amgen, 7/2025)
trastuzumab-dkst (Ogivri)	IV: Dose in 250 mL NS	24 h Fridge	Direct diluent into cake to reconstitute; swirl vial gently, allow to sit for 5 min. Gently invert bag to mix. Do not shake.			150 mg 420 mg	150 mg: 7.4 mL SWFI 420 mg: 20 mL BACT SW	150 mg=24 h 420 mg=28 d (w/Bact SW)	I (Mylan, 11/2024)
trastuzumab-qyyp (Trazimera)	IV: Dose in 250 mL NS	24 h Fridge	Direct diluent into cake to reconstitute; swirl vial gently, allow to sit for 5 min. Gently invert bag to mix. Do not shake.			150 mg 420 mg	150 mg: 7.4 mL SWFI 420 mg: 20 mL BACT SW	150 mg=24 h 420 mg=28 d (w/Bact SW)	I (Pfizer, 9/2025)
tremelimumab-actl (Injudo)	IV: 0.1 - 10 mg/mL NS or D5W	24 h	Gently invert bag to mix. Do not shake.	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	25 mg/1.25 mL 300 mg/15 mL	n/a	discard unused	I (AstraZeneca, 7/2024)
ublituximab-xiyy (Briumvi)	IV: Dose in total volume 250 mL NS	8 h	Gently invert bag to mix. Do not shake. If not used immediately, may store 24 h Fridge			150 mg/6 mL	n/a	discard unused	I (TG Therapeutics, 8/2025)
ustekinumab (Stelara)	IV: Dose in total volume 250 mL NS	8 h	Withdraw amount of fluid from bag equal to dose being added, before adding dose. Gently invert bag to mix. Do not shake.	Low sorbing tubing + 0.2 micron filter (11532269)	Filter	130 mg/26 mL	n/a	discard unused	I (Janssen, 4/2025)
vedolizumab (Entyvio)	IV: Dose in 250mL NS	12 h	Use a 21 to 25 ga needle; direct stream of SWFI to side wall of vial; gently swirl 15 seconds; let sit; can take up to 20 min. Once dissolved, gently invert vial 3 times. Gently invert bag to mix. Do not shake. If not used immediately, may store 24 h Fridge.			300 mg	300 mg: 4.8 mL SWFI	8 h	I (Takeda, 12/2025)
zolbetuximab (Vyloy)	IV: 5 mg/mL in NS	6 h	Direct diluent to vial wall to reconstitute; swirl vial gently, do not shake. Gently invert bag to mix. Do not shake. If not used immediately, may store 16 h Fridge.	LPB 0.2 micron filter (20027E)	Filter	100 mg 300 mg	100 mg: 5 mL SWFI 300 mg: 15 mL SWFI	discard unused	I (Astellas, 6/2025)

NOTE: BUD times reflect admixture in an ISO 5 LAFW within an ISO 7 environment. SCA compounding reduces BUD to 12 h room temp, 24 h fridge unless stability of product is less.

Approved by USP Standards, 2/2026