

January 2024 to December 2024 Antibiogram (% Susceptible) All Isolates (Inpatient and Outpatient)	Isolates	Amoxicillin + Clavulanate	Ampicillin	Ampicillin + Sulbactam	Aztreonam	Cefazolin	Cefepime	Cefotaxime	Cefoxitin	Ceftazidime	Ceftioxone	Ciprofloxacin	Clindamycin	Erythromycin	Gentamicin	Levofloxacin	Linezolid	Meropenem	Nitrofurantoin	Oxacillin	Penicillin	Piperacillin + Tazobactam	Trimethoprim + Sulfamethoxazole	Tetracycline	Tobramycin	Vancomycin
Gram Negative																										
Acinetobacter baumannii	6**	—	0*	66*	100*	—	100*	—	—	100*	0*	—	—	—	100*	—	—	100*	—	—	—	—	60*	—	100*	—
Citrobacter koseri	31	—	0	100	100	100	100	—	—	100	100	100	—	—	100	100	—	100	87*	—	—	100	100	—	100	—
Enterobacter cloacae	24**	—	4*	8*	95*	0*	95*	—	—	95*	78*	95*	—	—	100*	95*	—	100*	36*	—	—	100*	95*	—	95*	—
Escherichia coli	1002	—	48	53	99	90	99	9*	100*	98	98	79	—	—	90	83	—	100	98	—	—	99	71	—	92	—
Escherichia coli ESBL	76	—	0	39	0	0	0	4	—	1	0	25	—	—	78	44	—	100	93	—	—	97	38	—	73	—
Klebsiella oxytoca	11**	—	9*	72*	100*	63*	100*	—	—	100*	100*	83*	—	—	100*	83*	—	100*	100*	—	—	100*	77*	—	100*	—
Klebsiella pneumoniae	126	—	1	74	100	95	100	—	—	100	100	96	—	—	100	97	—	100	54	—	—	98	96	—	100	—
Morganella morganii	5**	—	0*	0*	60*	0*	100*	—	—	—	100*	75*	—	—	100*	75*	—	100*	0*	—	—	100*	100*	—	100*	—
Proteus mirabilis	83	—	89	92	100	92	100	—	—	100	98	96	—	—	97	98	—	100	0	—	—	100	95	—	97	—
Pseudomonas aeruginosa	61	—	—	0*	95	—	100	—	—	98	—	96	—	—	91	96	—	96	—	—	—	100	0*	—	100	—
Serratia marcescens	20**	—	0*	5*	45*	0*	100*	—	—	30*	52*	100*	—	—	100*	100*	—	100*	0*	—	—	75*	100*	—	85*	—
Stenotrophomonas maltophilia	4**	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	100*	—	—	—
Gram Positive																										
Enterococcus faecalis	174	—	99	—	—	—	—	—	—	—	—	—	—	16*	89	—	100	—	100	—	99	—	—	29	—	100
Enterococcus faecium	4**	—	75*	—	—	—	—	—	—	—	—	—	0*	100*	100*	—	100*	—	100*	—	50*	—	—	33*	—	100*
Streptococcus group B	102	100*	100	—	—	—	100	100	—	—	100	—	24*	18*	—	84	—	100*	—	—	100	—	100*	0*	—	100
MRSA	77	3	0	7	—	3	—	—	—	—	—	0*	73	16	100*	0*	100	—	100*	0	0	—	100	96	—	100
MSSA	82	100	—	100	—	100	—	—	—	—	—	—	77	57	—	—	100	—	100*	100	—	—	100	93	—	100
Staphylococcus epidermidis	25**	100*	—	100*	—	100*	—	—	—	—	—	—	53*	15*	—	—	100*	—	100*	100*	—	—	81*	77*	—	100*
Staphylococcus lugdunensis	11**	90*	—	90*	—	90*	—	—	—	—	—	—	88*	90*	—	—	100*	—	100*	90*	—	—	100*	90*	—	100*
Staphylococcus saprophyticus	52	100*	—	100*	—	100*	—	—	—	—	—	—	100*	0*	—	—	96	—	100	0*	—	—	96	100*	—	100*

** Inclusion of 30 isolates recommended for statistical significance by CLSI (Clinical and Laboratory Standards Institute).

* Indicates < 30 isolates tested for antibiotic listed

During 2024 zero Vancomycin Resistant Enterococcus faecalis and Vancomycin Resisant Enterococcus faecium were reported.

Greater than 50% of S. aureus isolates in the community are MRSA. These isolates are sensitive to Vancomycin, Linezolid, Daptomycin, and usually Trimethoprim / Sulfamethoxazole.

Therefore, critically ill patients with presumptive S. aureus should receive antimicrobial agents that treat MRSA until culture and sensitivity results are known.

Group B Streptococcus resistance to Clindamycin was not uniformly tabulated for antibiogram purposes in isolates from adults in 2023.

Abbreviations: MSSA = Methicillin Sensitive Staphylococcus aureus, MRSA = Methicillin Resistant Staphylococcus aureus